

# Biological Properties of *Thuja Orientalis* Linn

Priya Srivastava<sup>1</sup>, P. Kumar<sup>2</sup>, D. K. Singh<sup>2</sup>, V. K. Singh<sup>2,\*</sup>

<sup>1</sup>Department of Zoology, St. Xavier's College, Ranchi, Jharkhand, 834001, India

<sup>2</sup>Malacology Laboratory, Department of Zoology, D.D.U. Gorakhpur University, Gorakhpur, 273009, India

**Abstract** *Thuja orientalis* (Commonly- Morpankhi, Family- Cupressaceae) is an evergreen, monoecious trees or shrubs used in various forms of traditional medicines and homeopathy in various ways. In traditional practices *Thuja* is used for treatment of bronchial catarrh, enuresis, cystitis, psoriasis, uterine carcinomas, amenorrhoea and rheumatism. Recent researches in different parts of the world have shown that *T. orientalis* and its active component thujone have the great potential against a various health problems. *T. orientalis* preparations can be efficiently used against microbial/worm infection. It can be used as antioxidant, anticancer and anti-inflammatory agent. Instead of these effects, it can be also used as insecticidal, molluscicidal and nematocidal activity against different pests. The present review highlights the some important biological properties of *T. orientalis*.

**Keywords** *Thuja Orientalis*, Morpankhi, Thujone, Antimicrobial, Molluscicidal, Antimicrobial

## 1. Introduction

*Thuja orientalis* (Commonly- Morpankhi, Family- Cupressaceae) is a genus of coniferous trees. *T. orientalis* is an evergreen, monoecious trees or shrubs growing to 10-60 feet tall. The shoot are flat, leaves are scale like. The leaves are arranged in flattened fan shaped growing with resin glands [1]. Their leaves contain essential oils used to treat fungus infections, cancer, moles and parasitic worms. The essential oil derived from the leaves is toxic.  $\alpha$ -thujone is useful as an insecticide and an antihelminthic agent for the treatment of parasitic worms[2]. However,  $\alpha$ -thujone is a toxic substance that disrupts neurological signals in the brain. Ingestion of the essential oils of *Thuja* leaves can cause death[2]. Seed with a pair of narrow lateral wings, seedlings produce 2 cotyledons. The wood is light, soft and aromatic. *Thuja* poles also often used to make fence posts and rails. The wood of *Thuja* is commonly used for guitar sound boards[3]. It is used as a medicinal plant in various forms of traditional medicines like folk medicine, homeopathy and treatment of bronchial catarrh, enuresis, cystitis, psoriasis, uterine carcinomas, amenorrhoea and rheumatism[4-6]. Oil of *Thuja* contains thujone which has been studied for its GABA (gamma-aminobutyric acid) receptor antagonistic, with potentially lethal properties[2]. A yellow dye is obtained from the young branches[7]. *Thuja* is also occasionally used for treating diseases of skin, blood, gastrointestinal tract, kidney, brain, warty excrescences, spongy tumors[6]. Dubey and Batra[8,9] reported that the hepato-protective activities

and antioxidant activity of *Thuja occidentalis*. Anti-proliferative and apoptosis-inducing properties of *Thuja occidentalis* has been evaluated by Biswas *et al.*, [6].

## 2. Chemical Constituents

*Thuja orientalis* leaves contain rhodoxanthin, amentoflavone, hinokiflavone, quercetin, myricetin, carotene, xanthophylls and ascorbic acid. The fruit and roots are strongly aromatic. Distillation of the dried roots yields an essential oil having the following properties- Sp.gr.20<sup>0</sup>, 0.971[ $\alpha$ ]D, -22.5<sup>0</sup> nD<sup>20</sup>, 1.5055: acid val, 2.1 ester. Val. 26.27; ester. Val. After acetylation, 89.39; Carbonyls (as C<sub>10</sub>H<sub>16</sub>O), 5.65% and 50% in 7-8 vols of 95% alcohol[10]. The composition of the oil is as follows: a new bicyclic sesquiterpene 51.10; l-borneol, 17.10; bornyl acetate, 9.1;  $\alpha$ -thujone and camphor, 5.6; and a new sesquiterpenic alcohol. The seed yields fatty oil having the following composition palmitic 5.28, stearic, 7.3; C<sub>18</sub> unsaturated acids, 1829 (linolenic, 44.6%); and C<sub>20</sub> unsaturated acids, 6.10%. The heartwood contains aromadendrin, taxifolin, widdrene, cedrol, thujopsadiene, dehydro- $\alpha$ -curcumene,  $\beta$ -isobiotol and Curcumenether. It also contains an essential oil C is a complex blend of: Sesquiterpene hydrocarbons (cuparenes) 40; alcohols (Cedrol, widdrol, cuparenols) 50; monoterpenic acids[10]. Nickavar *et al.*, [11] 19 and 28 compounds have been identified in the volatile oils of the fruit and leaf, respectively, while the fruit oil contained  $\alpha$ -pinene (52.4%), 3-carene (14.2%),  $\alpha$ -cedrol (6.5%) and phellandrene (5.1%), the leaf oil contained  $\alpha$ -pinene (21.9%),  $\alpha$ -cedrol (20.3%), 3-carene (10.5%) and limonene (7.2%) as the main components.


Thujone is a ketone and a monoterpene that occurs naturally in two diastereomeric forms:  $\alpha$ -thujone and  $\beta$ -thujone[12].

\* Corresponding author:

vinaygkpuniv@gmail.com (Vinay Kumar Singh)

Published online at <http://journal.sapub.org/als>

Copyright © 2012 Scientific & Academic Publishing. All Rights Reserved



**Schema 1.** Structure of diastereomeric isomer of thujone

### 3. Biological Effects

The main constituents of essential oils mono- and sesquiterpenes including carbohydrates, phenols, alcohols, ethers, aldehydes and ketones are responsible for the biological activity of aromatic and medicinal plants[12,13]. *Thuja orientalis* is used internally in the treatment of coughs, haemorrhages, excessive menstruation, bronchitis, asthma, skin infections, mumps, bacterial dysentery, arthritic pains and premature blandness[14]. The leaves are antipyretic, astringent, diuretic, emmenagogue, emollient, expectorant, refrigerant, and stomachic[15,16]. Their use is said to improve the growth of hair[15]. The seed is aperients, lenitive and sedative. It is used internally in the treatment of palpitations, insomnia, nervous disorder and constipation in the elderly. The bark is used in the treatment of burn and scalds. The stems are used in the treatment of coughs, cold, dysentery, rheumatism and parasitic skin diseases[16]. Thujone was a weak inhibitor of acyl-CoA: lysophosphatidylcholine acyltransferase activity in mouse brain synaptosomes compared to psychoactive cannabinoids[17].

#### 3.1. Antibacterial Activity

Plants and their essential oils are potentially useful sources of antimicrobial compounds. *Thuja orientalis* contain large amounts of three substances (alpha, beta and gamma thujaplicin) that in low concentration would serve as chelators for *Solmonella typhimurium*[18]. *T. orientalis* was very effective in inhibiting the growth of serotypes c and d of *Solmonella mutans* (MIC less than or equal to 2.0-7.8 mg/ml)[19].

#### 3.2. Antifungal Activity

The essential oil showed antifungal activity in the inhibition zone against *Alternaria alternata* and *Currularia lunata* in a direct bioautography assay by lipophilic leaf extract of *T. orientalis*. Best bioactive component ( $R_f = 0.80$ ) were observed and noted for antifungal activity. It produced an in-

hibition zone of 30 and 22 mm in diameter against *A. alternata* and *C. Lunata*, respectively[20,21]. Mishra *et al.*[22] reported antifungal activity of aqueous leaf extract of *T. orientalis* against *Curvularia lunata*. The essential oils from leaves, twigs and stems of large trees and shrub-like trees of *Thuja sutchuenensis* were extracted by hydrodistillation and supercritical fluid extraction and analysed by GC and GC-MS. The essential oils exhibited a certain degree of antifungal activity against six strains of human pathogenic fungi[23].

#### 3.3. Antiviral Activity

The chemical composition of the essential oil of *T. orientalis* was determined by GC/MS analysis. Essential oils have been evaluated for their inhibitory activity against Sever Acute Respiratory Syndrome *Coronavirus* (SARS-Coronavirus) and Herpes Simplex Virus Type-1 (HSV-1) replication *in vitro* by visually scoring of the virus-induced cytopathogenic effect post-infection[24,25]. Several researches have demonstrated that allopathic extracts of *T. orientalis* could be used as strong antiviral agents against plant and animal viruses[26,27].

#### 3.4. Inflammatory Activity

Vascular inflammation is involved in the inhibition and progression of cardiovascular disease including atherosclerosis. Anti-vascular inflammatory activity of an aqueous extract of *T. orientalis* (ATO) and its possible mechanisms were investigated in human umbilical vein endothelial cells (HUVECs)[28]. Pre- incubation of ATO inhibited tumor necrosis factor and also inhibited U937 monocytes adhesion to HUVECs stimulated by tumor necrosis factor (TNF) suggesting that it may inhibit the binding of monocytes to endothelium. Furthermore, ATO significantly inhibited TNF-induced production of intracellular reactive oxygen species (ROS). Overall, ATO has an anti- inflammatory activity which is at least in part, is due to the decrease in the TNF-induced endothelial adhesion to monocytes by inhibiting intracellular ROS production, NF- $\kappa$ B activation and cell adhesion molecule in HUVECs[28,29].

#### 3.5. Anticancer Activity

Strong 5 $\alpha$ -reductase inhibitor are extracted and fractionated from *T. orientalis* and purified as diterpenes in isolated form[30]. The inhibitors are used either on their own or as active ingredients of therapeutics in the treatment of diseases caused by the over activity of 5 $\alpha$ -reductase or the hypersecretion of androgens, such as male baldness, androgenetic alopecia, hirsutism, acne, prostatomegaly and cancer of the prostate[30,31]. Dubey and Batra[8,9] reported that the hepato-protective activities and antioxidant activity of *Thuja occidentalis* linn. Anti-proliferative and apoptosis-inducing properties of thujone-rich fraction (TRF) separated from *Thuja occidentalis*. Their possible anti-cancer potentials have been noted in the malignant melanoma cell line A375[6]. Sunila *et al.*,[32] concluded that a polysaccharide,

or long-chain sugar molecule, derived from *Thuja* leaves extract decreased the inflammation caused by cancer. It also prevented the cancer from metastasizing, or spreading throughout the body.

### 3.6. Larvicidal Activity

Larvicidal activities of *T. orientalis* oil against 4<sup>th</sup>-instar larvae of *Aedes aegypti* and *Culex pipiens pallens* has been observed by Ju-Hyun *et al.*[33]. Larvicidal activity of *T. orientalis* leaf oil was higher than those of stem, fruit, and seeds oils. Essential oils of leaves and fruits of *T. orientalis* at 400 ppm caused 100% and 71.6% mortalities against *A. aegypti*[33]. The larvicidal activity was observed from various age class (I-III) and found strong mortality in age class of II of *T. orientalis* against *Aedes aegypti* and *Culex pipiens pallens* larvae. Leaf part and age class II of *T. orientalis* has strong larvicidal activity against *Aedes aegypti* and *Culex pipiens pallens*. Leaf oil of *T. orientalis* shows natural larvicides against *Aedes aegypti* and *Culex pipiens pallens*[33].

### 3.7. Insecticidal Activity

Leaf extracts of *T. orientalis* shows a repellent activity against *Chilo partellus*. *T. orientalis* ether extract (68.63%), acetone extracts (67.51%) have sufficient repellent action [34]. Foliar application of semi-solid crude extract of *T. orientalis* on maize was very effective against *Chilo partellus*[35].

### 3.8. Nematicidal Activity

Ethanol extract of *Thuja orientalis* leaf concentrations (20,40,60, and 80%) at 3 time intervals caused mortality in egg juvenile of *Meloidogyne incognita*[36]. It revealed a linear relationship between the concentration of the plant extract and the number of eggs hatched. Mortality of juveniles was directly proportional to the concentration and time of plant extracts[36].

### 3.9. Molluscicidal Activity

Singh and Singh[37] reported that the ethanol extract of *T. orientalis* leaf (24 h LC<sub>50</sub>- 32.74 mg/l) and column purified fraction (24 h LC<sub>50</sub>- 29.25 mg/l) were potent molluscicide against *Lymnaea acuminata*. Thujone (24 h LC<sub>50</sub>- 08.09 mg/l) was identified as active molluscicidal component in *T. orientalis*. The molluscicidal activity of leaf/fruit of *Thuja orientalis* and their active components/column purified fraction with synergist Piperonyl butoxide (PB) and MGK-264 (ENT 8184) was studied in binary combination (1:5) against *L. acuminata*. Combination of *T. orientalis* leaf/thujone or fruit powder/ column extract of *T. orientalis* fruit with PB or MGK-264 indicate synergised the toxicity up to 189.02 times. Toxicity of binary combination was increased hundreds folds than their individual components indicating synergistic action[38]. Sub-lethal (40% and 80% of 24h LC<sub>50</sub>) *in vivo* treatments of column purified fraction of *Thuja orientalis* and their active molluscicidal component

thujone significantly inhibited the acetylcholinesterase (AChE), acid and alkaline phosphatase (ACP/ALP) activities in the nervous tissue of *Lymnaea acuminata* [39].

It can be concluded from the ongoing literature that *T. orientalis* has the great potential against a number of health problem viz. bacterial, fungal and worm infection. It has antioxidant, antiviral, insecticidal nematicidal and molluscicidal activity. Recently, it has shown carcinogenic property. It needs greater attention by the researchers to explore its full potential and efficient use in the human welfare.

---

## REFERENCES

- [1] Farjon, A. 2005. Monograph of Cupressaceae of Sciadopitys. Royal Botanic Gardens, Kew. ISBN 1-84246-068-4.
- [2] Hold, K. M., Sirisoma, N. S., Ikeda, T., Narahashi, T. and Casida, J. E. 2000. Alpha-thujone (the active component of absinthe): gamma-aminobutyric acid type A receptor modulation and metabolic detoxification". Proc. Natl. Acad. Sci., USA. 97(8): 3826-3831. doi:10.1073/pnas.070042397.
- [3] Bucur, V. 1995. Acoustics of wood. Boca Raton: CRC Press. pp298.
- [4] Shimada K. 1956. Contribution to anatomy of the central nervous system of the Japanese upon the vernal arbour vitae. Okajimas Folia Anatomica Japonica, 28 (1), 207-227.
- [5] Baran, D. 1991. Arbor vitae, a guarantee of health. Revista Medico-Chirurgicala a Societatii de Medici si Naturalisti din Lasi, 95(3-4): 347-349.
- [6] Biswas R., Mandal S.K., Dutta S., Bhattacharyya S.S., Boujedaini N. and Khuda-Bukhsh AR. 2011. Thujone-Rich Fraction of *Thuja occidentalis* Demonstrates Major Anti-Cancer Potentials: Evidences from In Vitro Studies on A375 Cells. Evidence-Based Complementary and Alternative Medicine, Volume 2011, Article ID 568148, 16 pages. doi:10.1093/ecam/nej042.
- [7] Grieve, M. 1984. A modern Herbal. Penguin. ISBN 0-14-046-440-9.
- [8] Dubey S.K. and Batra, A. 2008. Hepatoprotective activity from ethanol fraction of *Thuja occidentalis* Linn. Asian Journal of Research in Chemistry, 1:32-35.
- [9] Dubey S.K. and Batra, A. 2009. Antioxidant activity of *Thuja occidentalis* linn. Asian Journal of Pharmaceutical and Clinical Research, 2: 73-76.
- [10] Anonymous. 1995. The Wealth of India, Raw Materials Vol. IX. Publication and information directorate, CSIR, New Delhi, India.
- [11] Nickavar B. Amin G and Parhami S. 2003. Volatile constituents of the fruit and leaf oils of *Thuja orientalis* L. grown in Iran. Z. Naturforsch. 58c, 171D172.
- [12] Perry, N. B., Anderson, R. E., Brennan, N. J., Douglas, M. H., Heaney, A. J., McGimpsey, J. A. and Smallfield, B. M. 1999. "Essential Oils from Dalmatian Sage (*Salvia officinalis* L.): Variations among Individuals, Plant Parts, Seasons, and Sites". J Agric Food Chem 47(5): 2048-2054.

- [13] Sokovic M., Glamoclija J., Marin P.D., Brkic D., and van Griensven L.J.D. 2010. Antibacterial effects of the essential oils of commonly consumed medicinal herbs using an in vitro model. *Molecules* 15: 7532 - 7546; doi: 10.3390 / molecules15117532.
- [14] Bown, D. 1995. *Encyclopedia of Herbs and their Uses*. ISBN-0-7513-020-31.
- [15] Yeung, Him-Che, 1985. *Handbook of Chinese Herbs and Formulas*. Institute of Chinese medicine, Los Angeles.
- [16] Duke, J. A. and Ayensu (1985). *Medicinal plants of China*. Reference Publications Inc. Michigan.
- [17] Greenberg, J. H., Mellors, A. & McGowan, J. C. 1978. Molar volume relationships and the specific inhibition of a synaptosomal enzyme by psychoactive cannabinoids. *J. Med. Chem.*, 21: 1208.
- [18] Akers, H. A., Abrego, V. A. and Garland, E. 1980. Thujaplicins from *Thuja plicata* as iron transport agents for *Salmonella typhimurium*. *J. Bacteriol*, 141(1): 164-168.
- [19] Chen, C.P., Lin, C.C. and Namba, T., 1989. Screening of Taiwanese crude drugs for antimicrobial activity against *Streptococcus mutans*. *J. Ethnopharmacol.* 27, 285–295.
- [20] Guleria, S. and Kumar, A. 2006. Antifungal activity of some Himalayan medicinal plants using direct bioautography. *J. of Cell and Molecular Biology*, 5: 95-98.
- [21] Guleria, S., Kumar, A. and Tikua, A. K. 2007. Chemical Composition and Fungitoxic Activity of Essential Oil of *Thuja orientalis* L. Grown in the North- Western Himalaya. *Z. Naturforsch*, 63: 211-214.
- [22] Mishra, M., Malik, S. and Tiwari, S. N. 1992. Allelopathic effect of certain botanicals against six fungal pathogens of rice. *Proceedings of first National Symposium on Allelopathy in Agroecosystems (Agriculture and Forestry)*, 191-193.
- [23] Lie H., Wang Y. Su C., Liang F., Su W., Hui M., Shaw P., Luo Y. 2010. Chemical composition and antifungal activity of essential oils of *Thuja sutchuenensis*, a critically endangered species endemic to China. *Nat Prod Comm.* 5(10): 1673-1676.
- [24] Hassan, H. T., Drize, N. J., Sadovinkova, E.Yu., Gan, O. I., Gohla, S. and Neth, R. D. 1996. TPSg. An anti- human immunodeficiency virus (HIV-1) agent, isolated from the Cupressaceae *Thuja occidentalis* L. (Arborvitae) enhances in vivo hemopoietic progenitor cells recovery in sublethally irradiated mice. In: *Immunol. Lett.*, 50: 119-22.
- [25] Monica, R. L., Antoine, M. S., Rosa, T., Giancarlo, A. S., Francesco, M., Ilaria, L., Roberto, G., Jindrich, C. and Hans W. D. 2008. Phytochemical analysis and in vitro antiviral activities of the essential oils of seven Lebanon species. *Chem. Biodivers.*, 5: 461-470.
- [26] Gohla, S. H., Haubeck, H. D. and Neth, R. D. 1988. Mutagenic activity of high molecular polysaccharide fractions isolated from the Cupressaceae *Thuja occidentalis* L.I. Macrophage- dependent induction of CD-4 positive T-helper (Th+) lymphocytes. In *Leukemia.*, 2(8): 528-33.
- [27] Offergeld, R., Reinecker, C., Gumz, E., Schrum, S., Treiber, R., Neth, R. D. and Gohla, S. H. 1992. Mutagenic activity of high molecular polysaccharide fractions isolated from the Cupressaceae *Thuja occidentalis* L.(Arborvitae) enhanced cytokinene- production by thyapolsaccharide, g- fraction (TPSg). In: *Leukemia*, 3: 1895-1915.
- [28] Moon, M. K., Kang, D. G., Lee, Y. J., Kim, J. S. and Lee, H. S. 2008. Inhibitory activity of *Thuja orientalis* on TNF- induced vascular cell adhesion in HUVECs. *The FASEB Journal*, 22: 1128.20-1128.
- [29] Berger, A., Monnard, I., Baur, M., Charbonnet, C., Safonova, I. and Jomard, A. 2002. Epidermal anti-inflammatory properties of 5, 11, 14, 20:3: effect on mouse ear edema, PGE2 level in cultured keratinocytes, and PPAR activation. Doi: 10.1186/1476-511X-1-5. Licensee Bio. Med. Central Ltd.
- [30] US Patent 5773005 1998. Purified flavonoid and diterpene 5 $\alpha$ -reductase inhibitor from *Thuja orientalis* for androgen-related diseases.
- [31] Muhammad ,I., Mossa, J. S., Al Yahya, M. A., Ramadan, A. S. and El Seraly, F. L. 1995. Further antibacterial diterpenes from the bark and leaves of *Juniperus procera*. *Hochst. ex. Endl. Phytotherapy Research*, 9: 584-588.
- [32] Sunila E.S., Hamsa T.P. and Kuttan G. 2011. Effect of *Thuja occidentalis* and its polysaccharide on cell-mediated immune responses and cytokine levels of metastatic tumor-bearing animals. *Pharm Biol.* 49(10): 1965-1073.
- [33] Ju-Hyun, J., Sang- Hyun, L., Moo- Key, K., Hoi- seon, L. 2005. Larvicidal activity of *Chamaecyparis obusta* and *Thuja orientalis* leaf oils against two mosquito species. *Agric. Chemi. and Biotech.* 48 (1), 26-28.
- [34] Dwivedi, S. C. and Shekhawat, N. B. 2004. Repellent effect of some indigenous plant extract against *Trogoderma granarium* (Everts). *Asian J. Exp. Sci.*, 18: 47-51.
- [35] Anju, B. and Sharma, V. K. 1999. Relative toxicity and persistence of plant products against maize stem borer on maize. *Annals of Plant Protection Sciences*, 7(2): 144-149.
- [36] Cannayane, I. and Rajendran, G. 2002. Allelochemic action of certain plant extracts on eggs and juveniles of *Meloidogyn incognita*. *Current Nematology*, 13: 83-89.
- [37] Singh A. and Singh V.K. 2009. Molluscicidal activity of *Saraca asoca* and *Thuja orientalis* against the fresh water snail *Lymnaea acuminata*. *Veterinary Parasitology*, 164, 206-210, doi- 10.1016/j. vetpar.2009.05.008, UK.
- [38] Singh, A. Kumar, P., Singh D.K. and Singh V.K. 2010. Toxicity to binary combination of *Saraca asoca* and *Thuja orientalis* with synergist piperonyl butoxide and MGK-264 against the fresh water snail *Lymnaea acuminata*. *The Bioscan*, 5(1), 13-18.
- [39] Singh, A., Srivastava P., Kumar, P., Singh D.K. and Singh V.K. 2011. Inhibition of acetylcholinesterase and phosphatases by active molluscicidal components of *saraca asoca* and *thuja orientalis* in the nervous tissue of *Lymnaea acuminata*. 3rd International Conference on Climate Change, Forest Resource and Environment (ICCFRE), The Ecoscan, Special issue-1, 87-92.